

PATIENT INFORMATION

Name: _____ DOB: _____ Height: _____ Weight: _____
 Address: _____ Phone: _____ Allergies: _____
 City, State, Zip: _____ Email: _____

REFERRAL STATUS

New Referral Dose or Frequency Change Order Renewal

Is this the first dose? Yes No, date of last infusion: _____ Line type: PIV PICC Port Other

DIAGNOSIS AND ICD-10 CODE

Age related Osteoporosis without current pathological fracture ICD-10 Code: _____
 Age related Osteoporosis with current pathological fracture ICD-10 Code: _____
 Diagnosis: _____ ICD-10 Code: _____

REQUIRED DOCUMENTATION

This signed order form by the provider H&P and Clinical/Progress notes supporting primary diagnosis
 Patient demographics AND insurance information Labs and Tests supporting primary diagnosis
 Serum creatinine and serum calcium level within 30 days of infusion DEXA scan results and/or FRAX score
 Pregnancy test, as applicable

List Tried & Failed Therapies, including duration of treatment (please comment specifically on bisphosphonates):

1)
2)

PREMEDICATION ORDERS

Acetaminophen 650mg PO prior to infusion Other: _____
 Diphenhydramine 25mg PO prior to infusion Other: _____

MEDICATION ORDERS

Dosing Reclast 5 mg IV once yearly

RN to manage VAD per company protocol and administer ordered therapy per manufacturer guideline
 RN to access/start and deaccess/discontinue VAD as appropriate for therapy administration
 RN to flush and lock VAD/CVAD per company protocol

Other: _____

EMERGENCY MEDICATIONS

Administer the following medications as needed for infusion-related reactions per company protocol:

Adults (weight >40kg):

Diphenhydramine 25mg-50mg PO
 Diphenhydramine 25mg-50mg slow IV push over 2-5 mins
 Acetaminophen 325mg-650mg PO
 Methylprednisolone 125mg slow IV push over at least 5 mins as tolerated
 Epinephrine 0.3mg IM/SQ, may repeat x1
 Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

Pediatrics (weight <40kg): (may adjust with weight changes)

Diphenhydramine 25mg PO
 Diphenhydramine 25mg slow IV push over 2-5 mins
 Acetaminophen 325mg PO
 Methylprednisolone 40mg slow IV push over at least 5 mins as tolerated
 Epinephrine 0.15mg (<30kg) or 0.3mg (>30kg) IM/SQ, may repeat x1
 Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

Oxygen 1-6LPM continuous flow per nasal cannula or face mask, titrate to maintain SpO2 of 95-100% (AIC/AIS only)

PRESCRIBER INFORMATION

Prescriber Name: _____ NPI Number: _____
 Office Phone: _____ Office Fax: _____
 Prescriber Signature: _____ Date: _____