

PATIENT INFORMATION

Name: _____ DOB: _____ Height: _____ Weight: _____
 Address: _____ Phone: _____ Allergies: _____
 City, State, Zip: _____ Email: _____

REFERRAL STATUS

New Referral Dose or Frequency Change Order Renewal

Is this the first dose? Yes No, date of last infusion: _____ Line type: PIV PICC Port Other

DIAGNOSIS AND ICD-10 CODE

Crohn's Disease ICD-10 Code: _____ Psoriatic Arthritis ICD-10 Code: _____
 Ulcerative Colitis ICD-10 Code: _____ Diagnosis: _____ ICD-10 Code: _____
 Plaque Psoriasis ICD-10 Code: _____

REQUIRED DOCUMENTATION

This signed order form by the provider TB Test Results
 Patient demographics AND insurance information Labs and Tests supporting primary diagnosis
 H&P and Clinical/Progress notes supporting primary diagnosis

List Tried & Failed Therapies, including duration of treatment: 1) _____ 2) _____

MEDICATION ORDERS

Medication	Indication	Dose	Route	Frequency
<input type="checkbox"/> Skyrizi (Risankizumab-rzaa)	Plaque Psoriasis or Psoriatic Arthritis	<input type="checkbox"/> 150 mg	SubQ	Week 0, 4, and every 12 weeks thereafter
	Crohn's Disease	Induction: <input type="checkbox"/> 600mg infused over 60 minutes	IV	Week 0, 4, 8
		Maintenance: <input type="checkbox"/> 180mg <input type="checkbox"/> 360mg	SubQ	Week 12, then every 8 weeks thereafter
	Ulcerative Colitis	Induction: <input type="checkbox"/> 1200mg infused over 120 minutes	IV	Week 0, 4, 8
		Maintenance: <input type="checkbox"/> 180mg <input type="checkbox"/> 360mg	SubQ	Week 12, then every 8 weeks thereafter

Hepatotoxicity in treatment of Inflammatory Bowel Disease. Drug induced liver injury during induction has been reported. Monitor LFT's and bilirubin at baseline and during induction, up to at least 12 weeks of treatment. Monitor thereafter according to routine patient management.

Refills*: X 6 months X 1 Year Other: _____

*(if not indicated, order will expire 1 year from date signed)

RN to manage VAD per company protocol and administer ordered therapy per manufacturer guideline
 RN to access/start and deaccess/discontinue VAD as appropriate for therapy administration
 RN to flush and lock VAD/CVAD per company protocol

Other: _____

EMERGENCY MEDICATIONS

Administer the following medications as needed for infusion-related reactions per company protocol:

Adults (weight >40kg):

Diphenhydramine 25mg-50mg PO
 Diphenhydramine 25mg-50mg slow IV push over 2-5 mins
 Acetaminophen 325mg-650mg PO
 Methylprednisolone 125mg slow IV push over at least 5 mins as tolerated
 Epinephrine 0.3mg IM/SQ, may repeat x1
 Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

Pediatrics (weight <40kg): (may adjust with weight changes)

Diphenhydramine 25mg PO
 Diphenhydramine 25mg slow IV push over 2-5 mins
 Acetaminophen 325mg PO
 Methylprednisolone 40mg slow IV push over at least 5 mins as tolerated
 Epinephrine 0.15mg (<30kg) or 0.3mg (>30kg) IM/SQ, may repeat x1
 Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

Oxygen 1-6LPM continuous flow per nasal cannula or face mask, titrate to maintain SpO2 of 95-100% (AIC/AIS only)

PRESCRIBER INFORMATION

Prescriber Name: _____ NPI Number: _____
 Office Phone: _____ Office Fax: _____
 Prescriber Signature: _____ Date: _____