SPECIALTY INFUSION CENTER
An Access Infusion Care Company

CABOTEGRAVIR/RIPIVIRINE ORDERS (Cabenuva)

CENTERS AT SIC
An Access Infusion Care Managed Company

REFERRAL CHECKLIST
Please ensure the following information is included with your referral:
☐ Patient Demographics and Contact Information
☐ Patient Insurance Information
☐ Completed H&P and Progress Notes Including Medications and Allergies
☐ Recent Lab Results/Testing/Screenings Supporting the Diagnosis
☐ Completed and Signed Referral
☐ Applicable HIV testing (HIV1 RNA, HIV drug resistance testing)
☐ CBCwdif/CMP results
Including this information will minimize delays with prior authorizations and ensure timely treatment for your patient. Thank you!

©Specialty Infusion Center
Print, complete, and fax completed form to (855) 857-3834

SPECIALTY INFUSION CENTER
An Access Infusion Care Company

CABOTEGRAVIR/RIPIVIRINE ORDERS (Cabenuva)

CENTERS AT SIC
An Access Infusion Care Managed Company

PATIENT INFORMATION	PRESCRIBER INFORMATION
Patient Name:	Prescriber Name:
Address:	NPI:
City, State, Zip:	Address:
Cell: Email:	City, State, Zip:
Ht: Wt: kg Last 4 of SSN: DOB:	Phone: Fax:
Allergies:	Contact Person:

REFERRAL STATUS ☐ **New Referral** ☐ **Continuation Order** ☐ **Dose or Frequency Change**

First dose? ☐ Yes ☐ No - Date of last injection:

DIAGNOSIS

☐ Dx Code/description:

REQUIRED DOCUMENTATION ☐ **Signed order** ☐ **Demographics and insurance info** ☐ **H&P/labs/tests/progress notes supporting dx**

☐ HIV drug resistance testing	☐ Oral lead in start date: ☐ N/A
☐ HIV-1 RNA less than 50 copies per mL (attach results)	Prior Therapies:

PRESCRIBING INFORMATION

LAB ORDERS

Frequency: Every Labs: ☐ CMP ☐ CBC w diff ☐ Other:

MEDICATION

Medication	Strength	Dose/Directions	Quantity
✕ **Cabotegravir/ Rivipirine (Cabenuva)** If applicable, patient must be on oral lead in therapy for at least 28 days. Therapy should be continued until the day of first injection.	**Monthly Dose** 600mg/3ml-900mg/3ml OR 400mg/2mL-600mg/2ml **Every 2-month Dose** 600mg/3ml-900mg/3ml	**Monthly Dose** ☐ Induction:600mg/900mg x 1 dose ☐ Maintenance:400mg/600mg monthly Begin maintenance dosing 1 month after induction **Every-2-Month Dose** ☐ Induction:600/900mg monthly x 2 doses ☐ Maintenance:600mg/900mg every 2 months Begin maintenance dosing 2 months after induction Inject Intramuscularly (2 injections at **separate gluteal sites**)	☐ 1 Dosing kit ☐ 1 Dosing kit Refills: ☐ 1 Dosing kit Refills:1 ☐ 1 Dosing kit Refills:

If not specified, order will be valid for 1 year from the date of signature.

OBSERVATION

Observe the patient for at least 10 minutes after the injection.

NURSING INSTRUCTIONS/EMERGENCY MEDICATION

RN to manage VAD per company protocol and administer ordered therapy per manufacturer guidelines Post-Infusion:50ml0.9%NS to flush residual priming volume	PIV:Flush 3-10mL0.9%NS before/after PICC/Midline/Port:Flush 10mL0.9%NS before/after, then 3mLHeparin 10u/mL(PICC/Midline)/5mLHeparin 100u/mL(Port)final

Administer the following medications as needed for infusion-related reactions per company protocol:
Adults>40kg: Diphenhydramine25-50mgPO or IVP(2-5min); Acetaminophen325-650mgPO; Methylprednisolone125mgIVP(≥5min); Epinephrine0.3mgIM/SQ, may repeat×1; 0.9%NS500mLIV over 30-60 min, may repeat×1 for hypotension.
Pediatrics/Adults<40kg: Diphenhydramine25mgPO or IVP(2-5min); Acetaminophen325mgPO; Methylprednisolone40mgIVP(≥5min); Epinephrine0.15mg(<30kg)/0.3mg(>30kg)IM/SQ, may repeat×1; 0.9%NS500mLIV over 30-60min, may repeat×1 for hypotension.
Oxygen(AIC/AIS only):1-6L/min via NC/face mask; titrate to maintain $SpO_2$95-100%.

Prescriber Signature X: _____ **Date:** _____

☐ **Product Substitution Permitted** ☐ **Dispense as Written**

©Specialty Infusion Center
Print, complete, and fax completed form to (855) 857-3834