SPECIALTY INFUSION CENTER
An Access Infusion Care Company

TEPROTUMUMAB-TRBW (Tepezza)

CENTERS AT SIC
An Access Infusion Care Managed Company

REFERRAL CHECKLIST

Please ensure the following information is included with your referral:

- ☐ Patient Demographics and Contact Information
- ☐ Patient Insurance Information
- ☐ Completed H&P and Progress Notes Including Medications and Allergies
- ☐ Recent Lab Results/Testing/Screenings Supporting the Diagnosis
- ☐ CBC w/diff/CMP/TSH/T3/T4/A1C lab results
- ☐ Tried and failed treatments, including high dose steroids if not contraindicated
- ☐ Smoking Cessation Attestation: if patient is a smoker, please supply a provider attestation that smoking cessation discussion has occurred between patient and provider.
- ☐ Eye Exam: Prescriber must be (or working in consult with) an ophthalmologist. Patient must have active, moderate to severe TED with documentation of 1 or more of the following:
  -lid retraction of more than 2mm
  -moderate or severe soft tissue involvement
  -proptosis at least 3mm above normal values for race and sex
  -periodic or constant diplopia
- ☐ Clinical Activity Score (CAS) Report in the most severely affected eye (must be at least 4)
- ☐ Negative Pregnancy Test if applicable
- ☐ Completed and Signed Referral

Including this information will minimize delays with prior authorizations and ensure timely treatment for your patient.
Thank you!

©Specialty Infusion Center
Print, complete, and fax completed form to (855) 857-3834

SPECIALTY INFUSION CENTER
An Access Infusion Care Company

TEPROTUMUMAB-TRBW (Tepezza)

CENTERS AT SIC
An Access Infusion Care Managed Company

PATIENT INFORMATION	PRESCRIBER INFORMATION
Patient Name:	Prescriber Name:
Address:	NPI:
City, State, Zip:	Address:
Cell: Email:	City, State, Zip:
Ht: Wt: kg Last 4 of SSN: DOB:	Phone: Fax:
Allergies:	Contact Person:

REFERRAL STATUS ☐ New Referral ☐ Continuation Order ☐ Dose or Frequency Change	
First dose? ☐ Yes ☐ No - Date of last infusion:	Line type: ☐ PIV ☐ PICC ☐ Port ☐ Other

DIAGNOSIS

☐ Dx Code/description:

REQUIRED DOCUMENTATION ☐ Signed order ☐ Demographics and insurance info ☐ H&P, labs, tests, progress notes supporting dx	
☐ Negative pregnancy test or documentation of contraceptive use	☐ Confirmed Dx from an Ophthalmologist (include documentation)
• Prior Therapies:	

PRESCRIBING INFORMATION

LAB ORDERS

Frequency: Every Labs: ☐ CMP ☐ CBCwdif ☐ CRP ☐ Sed Rate ☐ Other:

PREMEDICATIONS (administer 30 minutes prior to infusion)

Acetaminophen: ☐ 650mgPO ☐ 1000mgPO	Methylprednisolone: ☐ 100mgIV ☐ 125mgIV ☐ mgIV
Diphenhydramine: ☐ 25mgPO ☐ 50mgPO ☐ 25mgIV ☐ 50mgIV	Other: ☐

MEDICATION

Medication	Strength	Dose/Directions	Quantity
☒ **Teprotumumab-tbrw (Tepezza)**	500mg/vial	☐ **Induction:** 10mg/kg x 1 dose ☐ **Maintenance:** 20mg/kg every 3 weeks x 7 doses Infuse in 100ml(<1800mg)/250ml(>1800mg) total volume 0.9%NS IV over 90 minutes (Infusion #1 and #2), then over 60 minutes for remaining infusions if tolerated	☒ # sufficient for weight-based dose Refills:

Use current documented weight; round calculated dose to nearest vial size +/- 10% of dose.
If not specified, order will be valid for 1 year from the date of signature.

NURSING INSTRUCTIONS/EMERGENCY MEDICATION

RN to manage VAD per company protocol and administer ordered therapy per manufacturer guidelines Post-Infusion:Use 50ml0.9%NS to flush residual priming volume	PIV:Flush 3-10mL0.9%NS before/after PICC/Midline/Port:Flush 10mL0.9%NS before/after, then 3mLHeparin 10u/mL(PICC/Midline)/ 5mLHeparin 100u/mL(Port) final

Administer the following medications as needed for infusion-related reactions per company protocol:
Adults>40kg: Diphenhydramine25-50mgPO or IVP(2-5min); Acetaminophen325-650mgPO; Methylprednisolone125mgIVP(≥5min); Epinephrine0.3mgIM/SQ, may repeat×1; 0.9%NS500mL IV over 30-60min, may repeat×1 for hypotension.
Pediatrics/Adults<40kg: Diphenhydramine25mgPO or IVP(2-5min); Acetaminophen325mgPO; Methylprednisolone40mgIVP(≥5min); Epinephrine0.15mg(<30kg)/0.3mg(>30kg)IM/SQ, may repeat×1; 0.9%NS500mL IV over 30-60min, may repeat×1 for hypotension.
Oxygen(AIC/AIS only):1-6L/min via NC/face mask; titrate to maintain SpO_2 95-100%.

Prescriber Signature X: _____ **Date:** _____

☐ **Product Substitution Permitted** ☐ **Dispense as Written**

©Specialty Infusion Center
Print, complete, and fax completed form to (855) 857-3834